

# From Glycobiology to Glycotechnology

by

\*1  
Yoshihisa Nishikawa and Tetsuo Takahashi \*2

(Received on Sep. 28, 2001)

## Abstract

Glycoproteins are biopolymers that contain one or more oligosaccharide chain covalently linked to a protein. During the past 15 years, it has been generally accepted that the asparagine-linked oligosaccharide of glycoprotein is crucial for the function of glycoprotein. Therefore in the biotechnology, many useful human glycoproteins [*ie.* erythropoietin, tissue plasminogen activator, interferon(  $\beta$  and  $\gamma$  ), *etc.*] are produced by transfecting each human corresponding gene to mammalian cells that have the machinery to synthesize asparagine-linked oligosaccharide. In the present paper, our recent studies on the biosynthesis of lipid-linked oligosaccharide which is the precursor of the asparagine-linked oligosaccharide of glycoprotein were described. We have succeeded in the cloning of human genes of the GDP-mannose dependent mannosyltransferases which are involved in the first half of biosynthesis of the lipid-linked oligosaccharide. This approach may enable us, not only to clarify the regulation of the biosynthesis of lipid-linked oligosaccharide, but also to apply for the enzymatic synthesis of functional oligosaccharide, and for the diagnosis of human congenital disorders of glycosylation (CDG).

**Keywords:** sugar, carbohydrate, biotechnology, glycotechnology, glycobiology, glycoconjugates, glycoprotein, asparagine-linked oligosaccharide, lipid intermediate, gene cloning, cDNA, glycosylation, glycosyltransferase, mannosyltransferase, congenital disorders of glycosylation

---

\*1 Professor, Department of Applied Biochemistry.

\*2 Associate Professor, Department of Applied Biochemistry.